

Size-biased branching population measures and the multi-type $x \log x$ condition

Peter Olofsson
Mathematics Department
Trinity University
San Antonio, Texas

April 27, 2009

Abstract

We investigate the $x \log x$ condition for a general (Crump–Mode–Jagers) multi-type branching process with a general type space by constructing a size-biased population measure that relates to the ordinary population measure via an intrinsic martingale W_t . Sufficiency of the $x \log x$ condition for a non-degenerate limit of W_t is proved and conditions for necessity are investigated.

GENERAL BRANCHING PROCESS; XLOGX CONDITION; IMMIGRATION; SIZE-BIASED MEASURE

AMS 2000 SUBJECT CLASSIFICATION: PRIMARY 60J80

1 Introduction

The $x \log x$ condition is a fundamental concept for supercritical Galton-Watson branching processes, being the necessary and sufficient condition for the process to grow as its mean. In a Galton-Watson process with offspring mean $m = E[X] > 1$, let Z_n be the number of individuals in the n th generation and let $W_n = Z_n/m^n$. Then W_n is a nonnegative martingale and hence $W_n \rightarrow W$ for some random variable W . The *Kesten-Stigum Theorem* is

Theorem 1.1 *If $E[X \log^+ X] < \infty$ then $E[W] = 1$; if $E[X \log^+ X] = \infty$ then $W = 0$ a.s.*

where $\log^+ x = \max(0, \log x)$. It can further be shown that $P(W = 0)$ must either be 0 or equal the extinction probability and hence $E[X \log^+ X] < \infty$ implies that $W > 0$ exactly on the set of nonextinction (see *e.g.* Athreya and Ney (1972)).

The analog for general single-type branching processes appears in Jagers and Nerman (1984) and a partial result (establishing sufficiency) for general multi-type branching processes in Jagers (1989). Lyons, Pemantle and Peres (1995) give a slick proof of the Kesten-Stigum theorem based on comparisons between the Galton-Watson measure and another measure, the *size-biased* Galton-Watson measure, on the space of progeny trees. In Olofsson (1998), these ideas were further developed to analyze general single-type branching processes and the current paper proceeds to general multi-type branching processes with a general type space. In addition to providing a new proof of a known result, size-biased processes also provide tools to further analyze necessity of the $x \log x$ condition.

A crucial concept for the Lyons-Pemantle-Peres (LPP) proof is that of size bias. If the offspring distribution is $\{p_0, p_1, \dots\}$ and has mean $m = E[X]$, the size-biased offspring distribution is defined as

$$\tilde{p}_k = \frac{k p_k}{m}$$

for $k = 0, 1, 2, \dots$ where we note in particular that $\tilde{p}_0 = 0$. A size-biased Galton-Watson tree is constructed in the following way: Let \tilde{X} denote a random variable that has the size-biased offspring distribution and let the ancestor v_0 have a number \tilde{X}_0 of children. Pick one of these at random, call her v_1 , give her a number \tilde{X}_1 of children, and give her siblings ordinary Galton-Watson descendant trees. Pick one of v_1 's children at random, call her v_2 and give her a number \tilde{X}_2 of children, give her sisters ordinary Galton-Watson descendant trees, and so on and so forth. With P_n denoting the ordinary Galton-Watson measure restricted to the n first generations, \tilde{P}_n denoting the measure that arises from the above construction, and $W_n = Z_n/m^n$, it can be shown that the relation

$$d\tilde{P}_n = W_n dP_n, \tag{1.1}$$

holds. Hence, it is the martingale W_n that size-biases the Galton-Watson process. The construction of \tilde{P} can also be viewed as describing a Galton-Watson process with immigration where the immigrants are the siblings of the individuals on the path (v_0, v_1, \dots) . Thus, the measure P is the ordinary Galton-Watson measure and the size-biased measure \tilde{P} is the measure of a Galton-Watson process with immigration where the i.i.d. immigration group sizes are distributed as $\tilde{X} - 1$. The relation between P and \tilde{P} on the space of family trees can now be explored using results for processes with immigration which gives the final key to the proof.

The general idea of using size-bias in branching processes appeared before LPP. One early example is Joffe and Waugh (1982), where size-biased Galton-Watson processes show up in the study of ancestral trees of randomly sampled individuals. This approach was further explored by Olofsson and Shaw (2002) with a view toward biological applications. An approach similar to LPP appeared in Waymire and Williams (1996), developed simultaneously with, and independently of LPP. Later applications and extensions of the powerful LPP method include Kurtz *et al.* (1997), Geiger (1999), Athreya (2000), Biggins and Kyprianou (2004), and Lambert (2007).

To make this paper self-contained, we give a short review of general multi-type branching processes and their $x \log x$ condition in the next section. As in the Galton-Watson case, branching processes with immigration are crucial in the proof; for that purpose we briefly discuss processes with immigration in Section 3 following Olofsson (1996). The size-biased measure on the space of population trees and its relation to the ordinary branching measure is investigated in Section 4, and in Section 5, sufficiency of the $x \log x$ condition is proved. Finally, in Section 6, we discuss various conditions for necessity.

2 The $x \log x$ Condition for General Branching Processes

In a general branching process, individuals are identified by descent. The ancestor is denoted by 0, the children of the ancestor by 1, 2, ... and so on, so that the individual $x = (x_1, \dots, x_n)$ is the x_n th child of the x_{n-1} th child of...of the x_1 th child of the ancestor. The set of all individuals can thus be described as

$$I = \bigcup_{n=0}^{\infty} N^n$$

At birth, each individual is assigned a *type* s chosen from the *type space* S , equipped with some appropriate σ -algebra \mathcal{S} . The type space can be quite general; usually it is required to be a complete, separable metric (i.e., Polish) space. The type s determines a probability measure $P_s(\cdot)$, the *life law*, on the *life space* Ω , equipped with some appropriate σ -algebra \mathcal{F} . The information provided by a life $\omega \in \Omega$ may differ from one application to another but it must at least give the *reproduction process* ξ on $S \times R_+$. This process gives the sequence of birth times and types of the children of an individual. More precisely, let $(\tau(k), \sigma(k))$ be random variables on Ω denoting the birth time (age of the mother) and type of the k th child, respectively, and define

$$\xi(A \times [0, t]) = \#\{k : \sigma(k) \in A, \tau(k) \leq t\}$$

for $A \in \mathcal{S}$ and $t \geq 0$. We let $\tau(k) \equiv \infty$ if fewer than k children are born. The *population space* is defined as Ω^I , an outcome ω^I of which gives the lives of all individuals, together with the σ -algebra \mathcal{F}^I . The set of probability kernels $\{P_s(\cdot), s \in S\}$ defines a probability measure on $(\Omega^I, \mathcal{F}^I)$, the *population measure* P_s , where the ancestor's type is s .

With each individual $x \in I$, we associate its type σ_x , its birth time τ_x , and its life ω_x where σ_x is inherited from the mother (a function of the mother's life) and ω_x is chosen according to the probability distribution $P_{\sigma_x}(\cdot)$ on (Ω, \mathcal{F}) . The birth time τ_x is defined recursively by letting the ancestor be born at time $\tau_0 = 0$ and if x is the k th child of its mother y , we let $\tau_x = \tau_y + \tau(k)$. Note that τ_x and τ_y denote absolute time whereas $\tau(k)$ is the mother's age at x 's birth.

An important entity is the *reproduction kernel*, defined by

$$\mu(s, dr \times dt) = E_s[\xi(dr \times dt)]$$

the expectation of $\xi(dr \times dt)$ when the mother is of type s . This kernel plays the role of $m = E[X]$ in the simple Galton-Watson process and determines the growth rate of the process as $e^{\alpha t}$ where α is called the Malthusian parameter. We assume throughout that the process is supercritical, meaning that $\alpha > 0$. The existence of such an α is not automatic; sufficient conditions may be found in Jagers (1989, 1992). For the rest of this section, we

leave out further technical details and assumptions, instead focusing on the main definitions and results. The details can be found in Jagers (1989, 1992) and we simply refer to a process that satisfies all the conditions needed as a *Malthusian* process.

Given μ and α , we define the kernel $\hat{\mu}$ as

$$\hat{\mu}(s, dr) = \int_0^\infty e^{-\alpha t} \mu(s, dr \times dt) \quad (2.1)$$

which, under certain conditions, has eigenmeasure π and eigenfunction h given by

$$\begin{aligned} \pi(dr) &= \int_S \hat{\mu}(s, dr) \pi(ds) \\ h(s) &= \int_S h(r) \hat{\mu}(s, dr) \end{aligned} \quad (2.2)$$

where both π and $hd\pi$ can be normed to probability measures. The measure π is called the *stable type distribution* and $h(s)$ is called the *reproductive value* of an individual of type s . The interpretation of π and h is that π is the limiting distribution of the type of an individual chosen at random from a population, and $h(s)$ is a measure of how reproductive the type s tends to be, in a certain average sense. Moreover, after suitable norming, it can be shown that $hd\pi$ is probability measure that is the limiting type distribution backward in the family tree from the randomly sampled individual mentioned above. The mean asymptotic age of a random child-bearing in this backward sense is denoted by β and satisfies

$$\beta = \int_{S \times S \times R_+} t e^{-\alpha t} h(r) \mu(s, dr \times dt) \pi(ds) < \infty \quad (2.3)$$

To count, or measure, the population, *random characteristics* are used. A random characteristic is a real-valued process χ where $\chi(a)$ gives the contribution to the population of an individual of age a . Thus χ is a process defined on the life space and by letting χ_x be the characteristic pertaining to the individual x , the χ -counted population is defined as

$$Z_t^\chi = \sum_{x \in I} \chi_x(t - \tau_x)$$

which is the sum of the contributions of all individuals at time t (when the individual x is of age $t - \tau_x$). The simplest example of a random characteristic is $\chi(a) = I_{R_+}(a)$, the indicator for being born, in which case Z_t^χ is simply the total number of individuals born up to time t .

To capture the asymptotics of Z_t^χ , the crucial entity is the *intrinsic martingale* W_t , introduced by Nerman (1981) for single-type processes and generalized to multi-type processes in Jagers (1989). For its definition, denote x 's mother by mx and let

$$\mathcal{I}_t = \{x : \tau_{mx} \leq t < \tau_x\} \quad (2.4)$$

the set of individuals whose mothers are born at, or before, time t but who themselves are not yet born at time t . The set \mathcal{I}_t , sometimes referred to as the ‘‘coming generation,’’ generalizes the concept of generation in the Galton-Watson process. Now let

$$W_t = \frac{1}{h(\sigma_0)} \sum_{x \in \mathcal{I}_t} e^{-\alpha \tau_x} h(\sigma_x) \quad (2.5)$$

the individuals in \mathcal{I}_t summed with time- and type-dependent weights, normed by the reproductive value of the ancestor. It can be shown that W_t is a martingale with respect to the σ -algebra \mathcal{F}_t generated by the lives of all individuals born before t , and that $E_s[W_t] = 1$ for all $s \in S$. Hence, W_t plays the role that $W_n = Z_n/m^n$ does in the Galton-Watson process and the limit of Z_t^χ turns out to involve the martingale limit $W = \lim_{t \rightarrow \infty} W_t$. The main convergence result is of the form

$$e^{-\alpha t} Z_t^\chi \rightarrow \frac{E_\pi[\hat{\chi}(\alpha)]}{\alpha\beta} h(s)W$$

P_s -almost surely for π -almost all $s \in S$, as $t \rightarrow \infty$. Here, $\sigma_0 = s$ is the type of the ancestor, $E_\pi[\cdot] = \int_S E_s[\cdot] \pi(ds)$, and $\hat{\chi}(\alpha)$ is the Laplace transform of $\chi(a)$ evaluated at the point α . As in the Galton-Watson case, the question is when the martingale limit W is non-degenerate. As $W_t \rightarrow W$ P_s -a.s. and $E_s[W_t] = 1$, L^1 -convergence with respect to P_s is equivalent to $E_s[W] = 1$ (Durrett (2005), p.258). Note that although it is the process Z_t^χ that is of interest and not W_t itself, the asymptotics are determined by W_t , one of many examples in probability of the usefulness of finding an embedded martingale.

We are ready to formulate the general $x \log x$ condition and the main convergence result. For the reproduction process ξ , define the transform

$$\bar{\xi} = \int_{S \times R_+} e^{-\alpha t} h(r) \xi(dr \times dt) \quad (2.6)$$

which plays the role of X in the Galton-Watson process (in fact, in that case $\bar{\xi} = X/m$). For future reference, let us also state an alternative representation of $\bar{\xi}$. Denote the sequence of birth times and types in the process ξ by $\tau(1), \sigma(1), \tau(2), \sigma(2), \dots$ and so on. Then

$$\bar{\xi} = \sum_{i=1}^{\infty} e^{-\alpha \tau(i)} h(\sigma(i)) \quad (2.7)$$

The $x \log x$ condition and convergence result are given in the following theorem from Jagers (1989).

Theorem 2.1 *Consider a general multi-type supercritical Malthusian branching process with*

$$E_{\pi}[\bar{\xi} \log^+ \bar{\xi}] < \infty$$

Then $E_s[W] = 1$ for π -almost all s , from which it follows that

$$e^{-\alpha t} Z_t^{\chi} \rightarrow \frac{E_{\pi}[\hat{\chi}(\alpha)]}{\alpha \beta} h(s) W$$

in $L^1(P_s)$ for π -almost all s .

3 Processes with Immigration

As mentioned in the Introduction, branching processes with immigration are crucial to our proof and in this section, we state the main result for such processes. Consider a general branching process where new individuals immigrate into the population according to some point process $\eta(dr \times dt)$ with points of occurrence and types $(\tau_1, \sigma_1), (\tau_2, \sigma_2), \dots$. The k th immigrant initiates a branching process according to the population measure P_{σ_k} . The immigration process has the transform

$$\bar{\eta} = \int_0^\infty e^{-\alpha t} h(r) \eta(dr \times dt) = \sum_{k=1}^\infty e^{-\alpha \tau_k} h(\sigma_k)$$

and it can be shown that the process W_t is now a submartingale rather than a martingale (which is intuitively clear because offspring of immigrants may be added to the set \mathcal{I}_t). The limit of W_t is therefore not automatically finite but needs a condition on the immigration process, established by the following lemma from Olofsson (1996).

Lemma 3.1 *If $\bar{\eta} < \infty$ a.s., then $W_t \rightarrow W$ a.s. as $t \rightarrow \infty$ where $W < \infty$ a.s.*

4 The size-biased population measure

Recall that the LPP size-biased Galton-Watson measure was constructed from the size-biased offspring distribution. General branching processes require a more general concept of size-bias. In a general process, the offspring random variable X is replaced by the reproduction process ξ , the size of which is properly measured by the transform $\bar{\xi}$ which leads to the following definition.

Definition 4.1 *The size-biased life law \tilde{P}_s is defined as*

$$\tilde{P}_s(d\omega) = \frac{\bar{\xi}(\omega)}{h(s)} P_s(d\omega)$$

From Jagers (1992) we know that the eigenfunction h is finite and strictly positive so \tilde{P} is well-defined. The following lemma follows immediately from the definition of \tilde{P}_s .

Lemma 4.2 *Let P_s and \tilde{P}_s be as above and denote the set of realizations of reproduction processes by Γ , equipped with a σ -algebra \mathcal{G} . Then*

(i) *For $A \in \mathcal{F}$,*

$$\tilde{P}_s(A) = \frac{E_s[\bar{\xi}; A]}{h(s)}$$

(ii) For every \mathcal{G} -measurable function $g : \Gamma \rightarrow R$,

$$\tilde{E}_s[g(\xi)] = \frac{E_s[\bar{\xi}g(\xi)]}{h(s)}$$

Note that \tilde{P}_s is indeed a probability measure for all $s \in S$ because

$$\tilde{P}_s(\Omega) = \frac{1}{h(s)} E_s[\bar{\xi}] = 1$$

where $E_s[\bar{\xi}] = h(s)$ follows from the definition of $\bar{\xi}$ in (2.6) together with (2.1) and (2.2). Also note that a size-biased reproduction process always contains points because

$$\tilde{P}_s(\xi(S \times R_+) = 0) = \frac{1}{h(s)} E_s[\bar{\xi}; \xi(S \times R_+) = 0] = 0$$

in analogy with the size-biased offspring distribution in a Galton-Watson process ($\xi(S \times R_+)$ is the total number of offspring of an individual).

To construct the size-biased population measure, let \tilde{P}_s^t and P_s^t denote the restrictions of the measures P_s and \tilde{P}_s to the σ -algebra \mathcal{F}_t . The goal is to construct a measure \tilde{P}_s on $(\Omega^I, \mathcal{F}^I)$ that is such that

$$\tilde{P}_s^t(d\omega^I) = W_t(\omega^I) P_s^t(d\omega^I)$$

for all t , where W_t is the intrinsic martingale defined in (2.5). This measure is the direct extension of the size-biased measures from Lyons *et al.* (1995) and Olofsson (1998). The construction also involves the set \mathcal{I}_t , defined in (2.4), whose individuals all have mothers that are born up to time t . Thus, the type and birth time of an individual in \mathcal{I}_t is measurable with respect to \mathcal{F}_t which implies that W_t is also measurable with respect to \mathcal{F}_t .

The construction of the size-biased population measure extends the construction in Olofsson (1998) as follows. Start with the ancestor, now called v_0 , and choose her life ω_0 according to the size-biased distribution $\tilde{P}_s(d\omega_0) = \frac{\xi_0}{h(s)} P_s(d\omega_0)$. Pick one of her children, born in the reproduction process ξ_0 ,

such that the i th child is chosen with probability $\frac{e^{-\alpha\tau_i} h(\sigma_i)}{\bar{\xi}_0}$. Call this child v_1 , let her start a population according to the size-biased population law \tilde{P}_{σ_i}

and give her sisters independent descendant trees, such that sister j initiates a branching process according to the regular population law P_{σ_j} . Continue in this way and define the measure \tilde{P}_s to be the joint distribution of the random tree and the random path (v_0, v_1, \dots) . We shall borrow a term from Athreya (2000) and refer to the path (v_0, v_1, \dots) of chosen individuals as the *spine*.

Now fix an individual x in the set \mathcal{I}_t , defined in (2.4) and consider the probability \tilde{P}_s constrained by the individual x being chosen to be in the spine. Specifically, if $S(x, t)$ denotes the event that the individual x in \mathcal{F}_t is chosen to be in the spine and $A \in \mathcal{F}^I$, we consider the measure $\tilde{P}_s(\cdot; x)$ defined by

$$\tilde{P}_s^t(A; x) = \tilde{P}_s^t(A \cap S(x, t)) \quad (4.1)$$

Denote by i the individual in the first generation from whom x stems, that is, $x = (i, y)$ for some y . Hence, if x is in the n th generation, it is of the form $x = (x_1, x_2, \dots, x_n)$ where $x_1 = i$ and we let $y = (x_2, \dots, x_n)$. In words, y is the same individual as x when i is viewed as the ancestor. Let $\omega^{(j)}$ denote the lives of all individuals when j is viewed as the ancestor to obtain

$$\tilde{P}_s^t(d\omega^I; x) = \frac{\bar{\xi}_0}{h(s)} P_s(d\omega_0) \cdot \frac{e^{-\alpha\tau_i} h(\sigma_i)}{\bar{\xi}_0} \cdot \tilde{P}_{\sigma_i}^{t-\tau_i}(d\omega^{(i)}; y) \cdot \prod_{j \neq i} P_{\sigma_j}^{t-\tau_j}(d\omega^{(j)}) \quad (4.2)$$

where the first factor describes the size-biased choice of life of the ancestor. The second factor is the probability that the individual i in the first generation is chosen to be in the spine and the third factor describes the size-biased probability measure of the process starting from i , constrained by the individual y being in the spine. The fourth factor, finally, describes the regular population measures stemming from the individuals in the first generation who are not chosen to be in the spine.

The following proposition states the desired relation between the size-biased measure \tilde{P}_s^t and the regular population measure P_s^t .

Proposition 4.3 *Let \tilde{P}_s^t and P_s^t be the restrictions of \tilde{P}_s and P_s to the σ -algebra \mathcal{F}_t and let W_t be as in (2.5). Then*

$$\frac{d\tilde{P}_s^t}{dP_s^t} = W_t$$

Proof. Let $\tilde{P}_s(\cdot; x)$ be as in (4.1). Then

$$\tilde{P}_s^t(d\omega^I; x) = \frac{e^{-\alpha\tau_x} h(\sigma_x)}{h(s)} P_s^t(d\omega^I)$$

Further, note that, for the regular population measure we have

$$\begin{aligned} P_s^t(d\omega^I) &= P_s(d\omega_0) \prod_{j=1}^{\xi_0(t)} P_{\sigma_j}^{t-\tau_j}(d\omega^{(j)}) \\ &= P_s(d\omega_0) P_{\sigma_i}^{t-\tau_i}(d\omega^{(i)}) \prod_{j \neq i} P_{\sigma_j}^{t-\tau_j}(d\omega^{(j)}) \\ &= \bar{\xi}_0 P_s(d\omega_0) \frac{h(\sigma_i)}{\bar{\xi}_0} \frac{1}{h(\sigma_i)} P_{\sigma_i}^{t-\tau_i}(d\omega^{(i)}) \prod_{j \neq i} P_{\sigma_j}^{t-\tau_j}(d\omega^{(j)}) \end{aligned}$$

Now let $\tau_y(i)$ and $\sigma_y(i)$ denote the birth time and type of the individual y when i is viewed as the ancestor. We then have $\tau_x = \tau_i + \tau_y(i)$ and $\sigma_x = \sigma_y(i)$.

Multiply $P_s^t(d\omega^I)$ by $\frac{e^{-\alpha\tau_x} h(\sigma_x)}{h(s)}$ to obtain

$$\begin{aligned} &\frac{e^{-\alpha\tau_x} h(\sigma_x)}{h(s)} P_s^t(d\omega^I) \\ &= \frac{\bar{\xi}_0}{h(s)} P_s(d\omega_0) \cdot \frac{e^{-\alpha\tau_i} h(\sigma_i)}{\bar{\xi}_0} \cdot \frac{e^{-\alpha\tau_y(i)} h(\sigma_y(i))}{h(\sigma_i)} \cdot P_{\sigma_i}^{t-\tau_i}(d\omega^{(i)}) \cdot \prod_{j \neq i} P_{\sigma_j}^{t-\tau_j}(d\omega^{(j)}) \\ &= \frac{\bar{\xi}_0}{h(s)} P_s(d\omega_0) \cdot \frac{e^{-\alpha\tau_i} h(\sigma_i)}{\bar{\xi}_0} \cdot \tilde{P}_{\sigma_i}^{t-\tau_i}(d\omega^{(i)}; y) \cdot \prod_{j \neq i} P_{\sigma_j}^{t-\tau_j}(d\omega^{(j)}) \\ &= \tilde{P}_s^t(d\omega^I; x) \end{aligned}$$

by (4.2). Finally, sum over $x \in \mathcal{I}_t$ to get

$$\begin{aligned} \tilde{P}_s^t(d\omega^I) &= \sum_{x \in \mathcal{I}_t} \frac{e^{-\alpha\tau_x} h(\sigma_x)}{h(s)} P_s^t(d\omega^I) \\ &= W_t P_s^t(d\omega^I) \end{aligned}$$

where ω^I is suppressed but understood as the argument of τ_x , σ_x , and W_t . ■

To summarize, restricted to the σ -algebra \mathcal{F}_t , the size-biased population measure \tilde{P}_s relates to the regular population measure P_s through the Radon-Nikodym derivative $W_t = \frac{d\tilde{P}_s^t}{dP_s^t}$, a relation that is the straight-forward extension from Olofsson (1998) which, in turn, is the straight-forward extension of the original LPP method. This result is also similar in nature to Proposition 1 in Athreya (2000) which deals with a different martingale.

The individuals v_0, v_1, \dots in the spine are of particular interest in our analysis. From now on, we will use $\sigma_0, \sigma_1, \dots$ to denote the types of the individuals in the spine. The interarrival times are denoted T_1, T_2, \dots , that is, T_k is the time between the appearances of the $(k-1)$ th and k th individual. In the single-type case treated in Olofsson (1998), the individuals in the spine have lives that are i.i.d. but the situation is now much different with dependence between consecutive individuals introduced via types. As we shall see later, this dependence is also what precludes the proof of necessity of the $x \log x$ condition to carry over from the single-type case.

We now state important properties of the sequences of types and interarrival times in two lemmas. The first lemma deals with the type sequence alone. For the rest of this section, we use the notation \tilde{P} rather than \tilde{P}_s since the conditional probabilities we consider do not depend upon the initial type.

Lemma 4.4 *The sequence of types $(\sigma_0, \sigma_1, \dots)$ in the spine is a homogeneous Markov chain with transition probabilities*

$$\tilde{P}(\sigma_{k+1} \in dr | \sigma_k = s) = \frac{h(r)}{h(s)} \hat{\mu}(s, dr)$$

and stationary distribution $\nu(ds) = h(s)\pi(ds)$.

Proof. In a generic reproduction process ξ , denote the birth time and type of the i th offspring by $\tau(i)$ and $\sigma(i)$, respectively. Note the difference between $\sigma(i)$ and σ_i , the latter being the type of the i th individual in the spine. The transition probabilities satisfy

$$\tilde{P}(\sigma_{k+1} \in dr | \sigma_k = s) = \sum_i \tilde{P}(\sigma_{k+1} \in dr, v_{k+1} = i | \sigma_k = s)$$

$$\begin{aligned}
&= \sum_i \tilde{E}_s \left[\frac{e^{-\alpha\tau(i)} h(\sigma(i))}{\bar{\xi}} \delta_{\sigma(i)}(dr) \right] \\
&= \frac{1}{h(s)} \sum_i E_s \left[e^{-\alpha\tau(i)} h(\sigma(i)) \delta_{\sigma(i)}(dr) \right] \\
&= \frac{1}{h(s)} \int_0^\infty E_s \left[e^{-\alpha t} h(r) \xi(dr \times dt) \right] \\
&= \frac{h(r)}{h(s)} \hat{\mu}(s, dr)
\end{aligned}$$

where we used Lemma 4.2 applied to the function

$$g(\xi) = \sum_i \frac{e^{-\alpha\tau_i} h(\sigma_i)}{\bar{\xi}}$$

Next, let $\nu(ds) = h(s)\pi(ds)$. As

$$\int_S \hat{\mu}(s, dr) \pi(ds) = \pi(dr)$$

we get

$$\int_{s \in S} \frac{h(r)}{h(s)} \hat{\mu}(s, dr) \nu(ds) = \nu(dr)$$

and thus the Markov chain of types in the spine has stationary distribution $\nu = h d\pi$. ■

The second lemma deals with the sequence of types and interarrival times of the individuals in the spine.

Lemma 4.5 *The sequence of types and interarrival times $(\sigma_0, T_1, \sigma_1, T_2, \dots)$ of the individuals in the spine constitutes a Markov renewal process with transition kernel*

$$\tilde{P}(T_{k+1} \in dt, \sigma_{k+1} \in dr | \sigma_k = s) = \frac{h(r)}{h(s)} e^{-\alpha t} \mu(s, dr \times dt)$$

and the expected value of T_k when $\sigma_0 \sim \nu$ is

$$\tilde{E}_\nu[T_k] = \beta < \infty$$

where β was defined in (2.3).

Proof. Similarly to the proof of Lemma 4.4, we get

$$\begin{aligned} & \tilde{P}(T_{k+1} \in dt, \sigma_{k+1} \in dr | \sigma_k = s) \\ &= \sum_i \tilde{P}(T_{k+1} \in dt, \sigma_{k+1} \in dr, v_{k+1} = i | \sigma_k = s) \\ &= \frac{1}{h(s)} E_s[e^{-\alpha t} h(r) \xi(dr \times dt)] \\ &= \frac{h(r)}{h(s)} e^{-\alpha t} \mu(s, dr \times dt) \end{aligned}$$

and the expected value of T_k when σ_0 is chosen according to the stationary distribution $\nu = hd\pi$ is

$$\begin{aligned} \tilde{E}_\nu[T_k] &= \int_{S \times [0, \infty)} t \frac{h(r)}{h(s)} e^{-\alpha t} \mu(s, dr \times dt) h(s) \pi(ds) \\ &= \int_{S \times [0, \infty)} t e^{-\alpha t} h(r) \mu(s, dr \times dt) \pi(ds) \\ &= \beta \end{aligned}$$

by (2.3). ■

There is an interesting connection between the size-biased measure and the *stable population measure* from Jagers (1992). The latter is an asymptotic probability measure that is centered around a randomly sampled individual as $t \rightarrow \infty$. In such a stable population, the randomly sampled individual is born in a point process that has the size-biased distribution, the asymptotic type distribution as time goes backwards through the individual's line of descent is $hd\pi$, and the asymptotic mean age at a random child-bearing is

β . The transition probabilities in this backward chain also involve $\hat{\mu}(s, dr)$ but have weights that are expressed in terms of π rather than h as we have in the size-biased measure where time goes forward. This relation becomes clearer in a finite-type Galton Watson process where π and h are simply the left and right eigenvectors of the mean reproduction matrix.

5 Sufficiency of the $x \log x$ condition

We are soon ready to prove the general $x \log x$ theorem, Theorem 2.1, the key to which is the relation between \tilde{P}_s and P_s . Recall that the two are related through W_t which is a martingale under P_s and a submartingale under \tilde{P}_s . The following lemma relates the limiting behavior of W_t under $P_\pi(\cdot) = \int_S P_s(\cdot)\pi(ds)$ to its limiting behavior under $\tilde{P}_\nu(\cdot) = \int_S \tilde{P}_s(\cdot)h(s)\pi(ds)$.

Lemma 5.1 *Let $W = \limsup_t W_t$. Then*

$$(i) \tilde{P}_\nu(W = \infty) = 0 \quad \Rightarrow \quad E_\pi[W] = 1$$

$$(ii) \tilde{P}_\nu(W = \infty) = 1 \quad \Rightarrow \quad E_\pi[W] = 0$$

Proof. By Durrett (2005), p.239

$$\tilde{P}_\nu(A) = \tilde{E}_\nu[W; A] + \tilde{P}_\nu(A \cap \{W = \infty\})$$

If $\tilde{P}_\nu(W = \infty) = 0$, we have $\tilde{P}_\nu(A) = \tilde{E}_\nu[W; A]$ and get

$$\tilde{P}_\nu(W = 0) = \tilde{E}_\nu[W; W = 0] = 0$$

and hence

$$\tilde{E}_\nu[W] = \tilde{E}_\nu[W : W > 0]$$

$$= \tilde{P}_\nu(W > 0) = 1$$

Moreover, as

$$1 = \tilde{E}_\nu[W] = \int_S \tilde{E}_s[W]\nu(ds)$$

and as Fatou's lemma implies that $\tilde{E}_s[W] \leq 1$ for all s , we must have $\tilde{E}_s[W] = 1$ for ν -almost all $s \in S$. As $\pi \ll \nu$, we also get $E_\pi[W] = 1$.

Next, suppose that $\tilde{P}_\nu(W = \infty) = 1$. As W is an a.s. finite martingale limit under P_s , for π -almost all $s \in S$, we have

$$P_\nu(W = \infty) = \int_S P_s(W = \infty)h(s)\pi(ds) = 0$$

Hence, the measures $\tilde{P}_\nu(\cdot)$ and $E_\nu[W; \cdot]$ are mutually singular and as $\tilde{P}_\nu(W > 0) > 0$, we get

$$E_\nu[W] = E_\nu[W; W > 0] = 0$$

which implies that $E_\pi[W] = 0$ as well, and the proof of the lemma is complete. \blacksquare

Another connection between expected values under P_π and \tilde{P}_ν , directly involving the $x \log x$ condition, is given by the following corollary to Lemma 4.2.

Corollary 5.2

$$E_\pi[\bar{\xi} \log^+ \bar{\xi}] = \tilde{E}_\nu[\log^+ \bar{\xi}]$$

Proof. Choose $g(\xi) = \log^+ \bar{\xi}$ in Lemma 4.2 to obtain

$$\begin{aligned} E_\pi[\bar{\xi} \log^+ \bar{\xi}] &= \int_S E_s[\bar{\xi} \log^+ \bar{\xi}] \pi(ds) \\ &= \int_S h(s) \tilde{E}_s[\log^+ \bar{\xi}] \pi(ds) \\ &= \tilde{E}_\nu[\log^+ \bar{\xi}] \end{aligned}$$

\blacksquare

The logical structure of the proof of Theorem 2.1 is as follows: Assume that $E_\pi[\bar{\xi} \log^+ \bar{\xi}] < \infty$. By Corollary 5.2, we then have $\tilde{E}_\nu[\log^+ \bar{\xi}] < \infty$. If we can show that this in turn implies that $W < \infty$ almost surely with respect

to \tilde{P}_ν , we can invoke Lemma 5.1(i) to conclude that $E_\pi[W] = 1$ after which the proof is more or less complete. The gap is filled by the next lemma that utilizes results for general branching processes with immigration. It has been mentioned that such processes play a vital role in the LPP method and in the current setting, we observe that the individuals off the spine constitute a general branching process with immigration, the immigrants being the siblings of the individuals v_1, v_2, \dots in the spine in an obvious extension of the Galton-Watson case. To describe the immigration process formally, let $I_{j,k}$ be the indicator of the event that v_{j-1} 's k th child is *not* chosen to become v_j , denote the immigration time of the j th immigrant by τ_j , and denote the birth time and type of the k th individual in ξ_j by $\tau_k(j)$ and $\sigma_k(j)$, respectively. The immigration process η is

$$\eta(ds \times dt) = \sum_{j,k} \delta_{\sigma_k(j)}(ds) \delta_{\tau_k(j)}(dt - \tau_j) I_{j,k}$$

which has

$$\bar{\eta} = \sum_{j,k} h(\sigma_k(j)) e^{-\alpha\tau_j} e^{-\alpha\tau_k(j)} I_{j,k} \quad (5.1)$$

We are now ready for the last lemma needed for the proof of Theorem 2.1.

Lemma 5.3 *Consider a general branching process with immigration process η as above. If $\tilde{E}_\nu[\log^+ \bar{\xi}] < \infty$, then $W = \lim_{t \rightarrow \infty} W_t$ exists and is finite \tilde{P}_ν -a.s.*

Proof. First note that

$$\tilde{E}_\nu[\log^+ \bar{\xi}] < \infty \Rightarrow \sum_n \tilde{P}_\nu(\log^+ \bar{\xi} > cn) < \infty \quad \text{for all } c > 0$$

Now consider the sequence $\bar{\xi}_0, \bar{\xi}_1, \dots$ for the individuals in the spine. Recalling that ν is the stationary distribution of the Markov chain of types in the spine, we conclude that $\tilde{P}_\nu(\log^+ \bar{\xi} > cn) = \tilde{P}_\nu(\log^+ \bar{\xi}_n > cn)$ for all n , which gives

$$\sum_n \tilde{P}_\nu(\log^+ \bar{\xi}_n > cn) < \infty$$

By the first Borel–Cantelli lemma, we get

$$\tilde{P}_\nu(\log^+ \bar{\xi}_n > cn \text{ i.o.}) = 0 \quad (5.2)$$

which, by (5.1) gives

$$\begin{aligned} \bar{\eta} &\leq \sum_{j,k} h(\sigma_k(j)) e^{-\alpha\tau_j} e^{-\alpha\tau_k(j)} \\ &= \sum_{j=1}^{\infty} e^{-\alpha\tau_j} \bar{\xi}_j < \infty \tilde{P}_\nu - \text{a.s.} \end{aligned}$$

since the τ_j are sums of the T_k which, being the regeneration times in a Markov renewal process, obey the strong law of large numbers (Alsmeyer (1994)), so that $\tau_j \sim \beta j$ almost surely as $j \rightarrow \infty$ (recall that $\tilde{E}_\nu[T_k] = \beta < \infty$). The last sum above is a.s. finite because of the subexponential growth of the $\bar{\xi}_n$ established in (5.2). By Lemma 3.1 we conclude that $\lim_t W_t$ exists and is finite \tilde{P}_ν -a.s. ■

We now have all the preliminaries in place to prove Theorem 2.1.

Proof of Theorem 2.1. As $E_\pi[\bar{\xi} \log^+ \bar{\xi}] = \tilde{E}_\nu[\log^+ \bar{\xi}] < \infty$, Lemma 5.3(i) implies that $\tilde{P}_\nu(W = \infty) = 0$ and Lemma 5.1(ii) gives $E_\pi[W] = 1$. Moreover, as

$$E_\pi[W] = \int_S E_s[W] \pi(ds)$$

and as Fatou's lemma implies that $E_s[W] \leq 1$ for all s , we must have $E_s[W] = 1$ for π -almost all $s \in S$. We are done. ■

6 Necessity of the $x \log x$ condition

For single-type processes, the condition of having finite $x \log x$ moment is both sufficient and necessary. Using the size-bias method, this can be established by using the first and second Borel–Cantelli lemmas, respectively. However, in the multi-type setting, the main result in Jagers (1989) establishes only sufficiency and it is currently not known whether necessity holds.

The method of size-biased branching processes provides a way of investigating necessity, and although the second Borel–Cantelli lemma can not be used due to dependence, more general versions can be employed. This section is explorative in nature and does not provide any definite solutions but outlines two different approaches to establish necessity under additional conditions through the conditional Borel–Cantelli lemma and the Kochen–Stone lemma, respectively.

By “the $x \log x$ condition,” we mean the condition that $E_\pi[\bar{\xi} \log^+ \bar{\xi}] < \infty$. Hence, to establish necessity we need to assume that $E_\pi[\bar{\xi} \log^+ \bar{\xi}] = \infty$ and show that this assumption implies that $E_\pi[W] = 0$, invoking Lemma 5.1(ii) in an intermediate step. The logical structure parallels that of the proof of sufficiency: If $E_\pi[\bar{\xi} \log^+ \bar{\xi}] = \infty$, Lemma 5.2 yields that $\tilde{E}_\nu[\log^+ \bar{\xi}] = \infty$ which implies that

$$\sum_n \tilde{P}_\nu(\log^+ \bar{\xi} > cn) = \infty$$

which, since ν is the stationary distribution for the types in the spine, further implies that

$$\sum_n \tilde{P}_\nu(\log^+ \bar{\xi}_n > cn) = \infty$$

In the proof of sufficiency, this sum was finite and the first Borel–Cantelli lemma could be invoked to conclude that $\tilde{P}_\nu(\log^+ \bar{\xi}_n > cn \text{ i.o.}) = 0$ which led to the rest of the proof. However, as the reproduction process ξ_{n+1} is chosen according to a probability distribution that depends on the type σ_n which is determined by the parent reproduction process ξ_n , we cannot assume that the $\bar{\xi}_n$ are independent. Hence, we cannot invoke the second Borel–Cantelli lemma to conclude that, almost surely, $\log^+ \bar{\xi}_n > cn$ i.o. This marks a difference from the single-type case treated in Olofsson (1998) where sufficiency and necessity were established using the first and second Borel–Cantelli lemmas, respectively. Below we establish necessity of the $x \log x$ condition under various additional conditions.

The conditional Borel–Cantelli lemma states that if $\{\mathcal{F}_n\}$ is a filtration and $\{A_n\}$ a sequence of events with $A_n \in \mathcal{F}_n$, then

$$\{A_n \text{ i.o.}\} = \left\{ \sum_{n=1}^{\infty} P(A_n | \mathcal{F}_{n-1}) = \infty \right\}$$

see Durrett (2005). For us, $A_n = \{\log^+ \bar{\xi}_n > cn\}$, the σ -algebra \mathcal{F}_{n-1} gives the type σ_n of the n th individual in the spine, and we get

$$\tilde{P}_s(\log^+ \bar{\xi}_n > cn | \mathcal{F}_{n-1}) = \tilde{P}_{\sigma_n}(\log^+ \bar{\xi} > cn)$$

and the question becomes under which conditions

$$\sum_{n=1}^{\infty} \tilde{P}_{\sigma_n}(\log^+ \bar{\xi} > cn) = \infty \quad \tilde{P}_\nu - \text{a.s.}$$

given that

$$\sum_{n=1}^{\infty} \tilde{P}_\nu(\log^+ \bar{\xi} > cn) = \infty$$

One more step is necessary in order to invoke part (ii) of Lemma 5.1, namely to argue that $\log^+ \bar{\xi}_n > cn$ i.o implies that $W = \limsup_t W_t = \infty$ \tilde{P}_ν -a.s. We state this as a lemma.

Lemma 6.1 *If $\tilde{P}_\nu(\log^+ \bar{\xi}_n > cn \text{ i.o}) = 1$, then $\tilde{P}_\nu(W = \infty) = 1$.*

Proof. Consider W_{τ_n} , the value of W_t at the time of the arrival of the n th immigrant v_n . All the children of this immigrant except the one chosen to become v_{n+1} belong to \mathcal{I}_{τ_n} , thus, these children form a subset of \mathcal{I}_{τ_n} . The k th child is born at time $\tau_n + \tau(k)$ and has type $\sigma(k)$ where $\tau(k)$ and $\sigma(k)$ are the points in the reproduction process ξ_n of v_n . Let I_k be the indicator of the event that the k th child of v_n is *not* chosen to become v_{n+1} to obtain

$$\begin{aligned} W_{\tau_n} &= \frac{1}{h(\sigma_0)} \sum_{x \in \mathcal{I}_{\tau_n}} h(\sigma_x) e^{-\alpha \tau_x} \\ &\geq \frac{e^{-\tau_n}}{h(\sigma_0)} \sum_{k=1}^{\infty} h(\sigma(k)) e^{-\alpha \tau(k)} I_k \\ &\geq \frac{e^{-\tau_n}}{h(\sigma_0)} (\bar{\xi}_n - C) \end{aligned}$$

where $C = \sup_s h(s) < \infty$ (Jagers (1989)) and we recall (2.7). Adjust the last part of the proof of Lemma 5.3 to conclude that $W = \limsup_t W_t = \infty$ \tilde{P}_ν -a.s. \blacksquare

Our first result establishes necessity of the $x \log x$ condition under the additional assumption that there is a type that is revisited infinitely often in the spine.

Proposition 6.2 *If the Markov chain of types $\sigma_0, \sigma_1, \dots$ in the spine has one positive recurrent state r such that $E_r[\bar{\xi} \log^+ \xi] = \infty$ and if, for π -almost all starting types s , there exists k such that $\sigma_k = r$ a.s., then the $x \log x$ condition is necessary.*

Note: The Markov chain is on a general state space but we here use “positive recurrence” in its elementary meaning, namely, that $\tilde{E}_r[T_r] < \infty$ where $T_r = \inf\{n > 0 : \sigma_n = r\}$.

Proof. The result follows from the following observation regarding infinite series. Let $a_n \geq 0$ be a decreasing sequence of real numbers such that $\sum_n a_n = \infty$, let X_1, X_2, \dots be i.i.d. nonnegative and integer-valued random variables with finite mean μ , and let $T_n = X_1 + X_2 + \dots + X_n$. Then

$$\sum_{n=1}^{\infty} a_{T_n} = \infty \quad \text{a.s.}$$

This holds because if $k \geq \mu$ is an integer, then $a_{T_n} \geq a_{nk}$ a.s. for large n by the strong law of large numbers and obviously $\sum_n a_{nk} = \infty$ for all fixed k . Finally, apply this result to $a_n = \tilde{P}_s(\log^+ \bar{\xi} > cn)$ with X_1, X_2, \dots being the consecutive inter-return times to the state s in the Markov chain of types in the spine (that is, the X_k are i.i.d. copies of T_r above). \blacksquare

One instance when the assumption of Proposition 6.2 follows from the $x \log x$ condition is if the type space is finite. Indeed, if there are n types, we have

$$\infty = E_\pi[\bar{\xi} \log^+ \xi] = \sum_{k=1}^n E_r[\bar{\xi} \log^+ \xi] \pi(r)$$

which implies that we must have $E_r[\bar{\xi} \log^+ \xi] = \infty$ for at least one r . In particular, this establishes necessity of the $x \log x$ condition for the ordinary

multi-type Galton–Watson process with a finite type space (under the usual assumptions of positive regularity and nonsingularity, see Athreya and Ney (1972)).

Another approach is to consider the rate of convergence of the type chain $\sigma_0, \sigma_1, \dots$ toward its stationary distribution ν . To simplify the analysis, let $Y = \lfloor \log^+ \bar{\xi} \rfloor$, the integer part of $\log^+ \bar{\xi}$, which has finite mean under \tilde{P}_ν if and only if $\log^+ \bar{\xi}$ does. We then have

Proposition 6.3 *Suppose that*

$$\sum_{n \geq 1} n \tilde{E}_\nu \left| \frac{1}{n} \sum_{k=1}^n \tilde{P}_{\sigma_k}(Y = n) - \tilde{P}_\nu(Y = n) \right| < \infty$$

Then the $x \log x$ condition is necessary.

Proof. The condition in the proposition implies that

$$\sum_{n \geq 1} n \left| \frac{1}{n} \sum_{k=1}^n \tilde{P}_{\sigma_k}(Y = n) - \tilde{P}_\nu(Y = n) \right| < \infty \quad \tilde{P}_\nu\text{-a.s.}$$

which yields

$$\begin{aligned} \sum_{k=1}^{\infty} \tilde{P}_{\sigma_k}(Y > k) &= \sum_{k=1}^{\infty} \sum_{n > k} \tilde{P}_{\sigma_k}(Y = n) \\ &= \sum_{n > 1} n \left(\frac{1}{n} \sum_{k=1}^n \tilde{P}_{\sigma_k}(Y = n) \right) \\ &\geq \sum_{n > 1} n \tilde{P}_\nu(Y = n) - \sum_{n \geq 1} n \left| \frac{1}{n} \sum_{k=1}^n \tilde{P}_{\sigma_k}(Y = n) - \tilde{P}_\nu(Y = n) \right| \\ &= \infty \end{aligned}$$

since the first term equals $\tilde{E}_\nu[Y]$ which is infinite if $\tilde{E}_\nu[\log^+ \bar{\xi}]$ is infinite, which is the case if the $x \log x$ condition does not hold. The second term is finite by assumption. \blacksquare

Note that if $\{\sigma_n\}$ is Harris recurrent (Alsmeyer (1994), Durrett (2005)) with stationary distribution ν , the ergodic theorem yields

$$\frac{1}{n} \sum_{k=1}^n \tilde{P}_{\sigma_k}(Y = j) \rightarrow \tilde{P}_\nu(Y = j) \quad \tilde{P}_\nu\text{-a.s.}$$

for all j so our condition means that this convergence is, in some vague sense, “fast enough.”

Another generalization of Borel–Cantelli is the Kochen–Stone lemma that states that if $\sum_n P(A_n) = \infty$, then

$$P(A_n \text{ i.o.}) \geq \limsup_n \frac{\{\sum_{k=1}^n P(A_k)\}^2}{\sum_{1 \leq j, k \leq n} P(A_j \cap A_k)}$$

We can apply it to prove

Proposition 6.4 *Let $A_n = \{\log^+ \bar{\xi}_n > cn\}$. If the (indicators of the) A_n are pairwise negatively correlated, the $x \log x$ condition is necessary.*

Proof. Because $\tilde{P}_\nu(A_j \cap A_k) \leq \tilde{P}_\nu(A_j)\tilde{P}_\nu(A_k)$, we get

$$\tilde{P}_\nu(A_n \text{ i.o.}) \geq \limsup_n \frac{\{\sum_{k=1}^n \tilde{P}_\nu(A_k)\}^2}{\sum_{1 \leq j, k \leq n} \tilde{P}_\nu(A_j \cap A_k)} \geq 1$$

■

7 Acknowledgements

The author would like to thank Gerold Alsmeyer and Olle Häggström for fruitful discussions related to the results in Section 6, the former of which took place more than 10 years ago.

8 References

- ALSMEYER, G. (1994) On the Markov renewal theorem. *Stoch. Proc. Appl.* **50**, 37–56.
- ATHREYA, K.B. (2000) Change of measures for Markov chains and the LlogL theorem for branching processes. *Bernoulli* **6(2)**, 323–338.

- ATHREYA, K.B. and NEY, P. (1972) *Branching Processes*. Springer, Berlin.
- BIGGINS, J.D. and KYPRIANOU, A. E. (2004) Measure change in multitype branching. *Adv. in Appl. Prob.* **36**(2), 544–581
- DURRETT, R. (2005) *Probability: Theory and Examples*, third edition. Duxbury Press.
- GEIGER, J. (1999) Elementary new proofs of classical limit theorems for Galton-Watson processes. *J. Appl. Prob* **36**, 301–309
- JAGERS, P. (1989) General branching processes as Markov fields. *Stoch. Proc. Appl.* **32**, 183–212.
- JAGERS, P. (1992) Stabilities and instabilities in population dynamics. *J. Appl. Prob.* **29**, 770–780.
- JAGERS, P. and NERMAN, O. (1984) The growth and composition of branching populations. *Adv. Appl. Prob.* **16**, 221–259.
- JOFFE, A. and WAUGH, W. A. O’N. (1982) Exact distributions of kin numbers in a Galton-Watson process. *J. Appl. Prob.* **19**(4), 767–775.
- KURTZ, T.G., LYONS, R., PEMANTLE, R. and PERES, Y. (1997) A conceptual proof of the Kesten-Stigum theorem for multitype branching processes. In K.B. Athreya and P. Jagers (eds), *Classical and Modern Branching Processes*, IMA Vol. Math. Appl. **84**, 181–186. New York: Springer-Verlag.
- LAMBERT, A. (2007) Quasi-stationary distributions and the continuous-state branching process conditioned to be never extinct. *Elec. J. Prob* **12**, 420–446
- LYONS, R., PEMANTLE, R. and PERES, Y. (1995) Conceptual proofs of $L \log L$ criteria for mean behavior of branching processes. *Annals of Probability* **3**, 1125–1138.
- NERMAN, O. (1981) On the convergence of supercritical general (C–M–J) branching processes. *Z. Wahrscheinlichkeitsth.* **57**, 365–395.

- OLOFSSON, P. (1996) General branching processes with immigration. *J. Appl. Prob.*, **33**, 940–948.
- OLOFSSON, P. (1998) The $x \log x$ condition for general branching processes. *J. Appl. Prob.*, **35**, 537–544.
- OLOFSSON, P. and SHAW (2002). Exact sampling formulas for multi-type Galton-Watson processes. *J. Math. Biol.*, **45**, 279–293
- WAYMIRE, E. and WILLIAMS, S.C. (1996) A cascade decomposition theory with applications to Markov and exchangeable cascades. *Trans. Amer. Math. Soc.* **348(2)**, 585–632.

Peter Olofsson
Mathematics Department
Trinity University
One Trinity Place
San Antonio, TX 78212

polofsso@trinity.edu

www.trinity.edu/polofsso