



PUTNAM SEMINAR
FALL 2018

QUIZ 2
DUE SEPTEMBER 5

Name: _____

Start Time: _____

End Time: _____

Problem 1. Let S be a subset of $\{1, 2, \dots, 2n\}$ with $n + 1$ elements. Show that one can choose distinct elements $a, b \in S$ such that a divides b .

Problem 2. For a partition π of $\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$, let $\pi(x)$ be the number of elements in the part containing x . Prove that for any two partitions π and π' , there are two distinct numbers x and y in $\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$ such that $\pi(x) = \pi(y)$ and $\pi'(x) = \pi'(y)$.

Problem 3. Let a_j, b_j, c_j be integers for $1 \leq j \leq N$. Assume for each j , at least one of a_j, b_j, c_j is odd. Show that there exist integers r, s, t such that $ra_j + sb_j + tc_j$ is odd for at least $4N/7$ values of j , $1 \leq j \leq N$.

Problem 4. Prove that, for every set $X = \{x_1, x_2, \dots, x_n\}$ of real numbers, there exists a non-empty subset S of X and an integer m such that

$$\left| m + \sum_{s \in S} s \right| \leq \frac{1}{n+1}$$

Problem 1. Let S be a subset of $\{1, 2, \dots, 2n\}$ with $n + 1$ elements. Show that one can choose distinct elements $a, b \in S$ such that a divides b .

Problem 2. For a partition π of $\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$, let $\pi(x)$ be the number of elements in the part containing x . Prove that for any two partitions π and π' , there are two distinct numbers x and y in $\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$ such that $\pi(x) = \pi(y)$ and $\pi'(x) = \pi'(y)$.

Problem 3. Let a_j, b_j, c_j be integers for $1 \leq j \leq N$. Assume for each j , at least one of a_j, b_j, c_j is odd. Show that there exist integers r, s, t such that $ra_j + sb_j + tc_j$ is odd for at least $4N/7$ values of j , $1 \leq j \leq N$.

Problem 4. Prove that, for every set $X = \{x_1, x_2, \dots, x_n\}$ of real numbers, there exists a non-empty subset S of X and an integer m such that

$$\left| m + \sum_{s \in S} s \right| \leq \frac{1}{n+1}$$