

# Periodic functions and Fourier series

Ryan C. Daileda



Trinity University

Partial Differential Equations  
February 4, 2014

**Goal:** Given a function  $f(x)$ , write it as a linear combination of cosines and sines, e.g.

$$\begin{aligned} f(x) &= a_0 + a_1 \cos(x) + a_2 \cos(2x) + \cdots + b_1 \sin(x) + b_2 \sin(2x) + \cdots \\ &= a_0 + \sum_{n=1}^{\infty} (a_n \cos(nx) + b_n \sin(nx)). \end{aligned}$$

### Important Questions:

1. Which  $f$  have such a *Fourier series expansion*?

Difficult to answer completely. We will give sufficient conditions only.

2. Given  $f$ , how can we determine  $a_0, a_1, a_2, \dots, b_1, b_2, \dots$ ?

We will give explicit formulae. These involve the ideas of *inner product* and *orthogonality*.

# Periodicity

**Definition:** A function  $f(x)$  is  $T$ -periodic if

$$f(x + T) = f(x) \text{ for all } x \in \mathbb{R}.$$

**Remarks:**


- If  $f(x)$  is  $T$ -periodic, then  $f(x + nT) = f(x)$  for any  $n \in \mathbb{Z}$ .
- The graph of a  $T$ -periodic function  $f(x)$  repeats every  $T$  units along the  $x$ -axis.
- To give a formula for a  $T$ -periodic function, state that “ $f(x) = \dots$  for  $x_0 \leq x < x_0 + T$ ” and then *either*:
  - \*  $f(x + T) = f(x)$  for all  $x$ ; OR
  - \*  $f(x) = f\left(x - T \left\lfloor \frac{x - x_0}{T} \right\rfloor\right)$  for all  $x$ .

# Examples

1.  $\sin(x)$  and  $\cos(x)$  are  $2\pi$ -periodic.
2.  $\tan(x)$  is  $\pi$ -periodic.
3. If  $f(x)$  is  $T$ -periodic, then:
  - $f(x)$  is also  $nT$ -periodic for any  $n \in \mathbb{Z}$ .
  - $f(kx)$  is  $T/k$ -periodic.
4. For  $n \in \mathbb{N}$ ,  $\cos(nkx)$  and  $\sin(nkx)$  are:
  - $2\pi/nk$ -periodic.
  - *simultaneously*  $2\pi/k$ -periodic.
5. If  $f(x)$  is  $T$ -periodic, then

$$\int_a^{a+T} f(x) dx = \int_0^T f(x) dx \text{ for all } a.$$

## 6. The 2-periodic function with graph




can be described by

$$f(x) = \begin{cases} x & \text{if } 0 < x \leq 2, \\ f(x+2) & \text{otherwise,} \end{cases}$$

or

$$f(x) = x - 2 \left[ \frac{x}{2} \right].$$

## 7. The 1-periodic function with graph



can be described by


$$f(x) = \begin{cases} 0 & \text{if } 0 < x \leq 1/2, \\ 2x - 1 & \text{if } 1/2 < x \leq 1, \\ f(x + 1) & \text{otherwise.} \end{cases}$$

# Piecewise smoothness

**Definition:** Given a function  $f(x)$  we define

$$f(c+) = \lim_{x \rightarrow c^+} f(x) \text{ and } f(c-) = \lim_{x \rightarrow c^-} f(x).$$

**Example:** For the following function we have:



$$f(0+) = 0,$$

$$f(0-) = -1,$$

$$f(1+) = f(1) = f(1-) = 1,$$

$$f(2+) = 2,$$

$$f(2-) = 1.$$


**Remark:**  $f(x)$  is continuous at  $c$  iff  $f(c) = f(c+) = f(c-)$ .

**Definition 1:** We say that  $f(x)$  is *piecewise continuous* if


- $f$  has only finitely many discontinuities in any interval, and
- $f(c+)$  and  $f(c-)$  exist for all  $c$  in the domain of  $f$ .

**Definition 2:** We say that  $f(x)$  is *piecewise smooth* if  $f$  and  $f'$  are both piecewise continuous.

Good:



Bad:



**Remark:** A piecewise smooth function *cannot* have: vertical asymptotes, vertical tangents, or “strange” discontinuities.



# Which functions have Fourier series?

We noted earlier that the functions

$$\cos(nx), \sin(nx), \quad (n \in \mathbb{N})$$

are all  $2\pi$ -periodic. It follows that if

$$f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos(nx) + b_n \sin(nx)),$$

then  $f(x)$  *must* also be  $2\pi$ -periodic.

However,  $2\pi$ -periodicity *does not* guarantee that  $f(x)$  has a Fourier series expansion.

But if we also require  $f(x)$  to be piecewise smooth...

# Existence of Fourier series

## Theorem

If  $f(x)$  is a piecewise smooth,  $2\pi$ -periodic function, then there are (unique) Fourier coefficients  $a_0, a_1, a_2, \dots$  and  $b_1, b_2, \dots$  so that


$$\frac{f(x+) + f(x-)}{2} = a_0 + \sum_{n=1}^{\infty} (a_n \cos(nx) + b_n \sin(nx))$$

for all  $x$ . This is called the Fourier series of  $f(x)$ .


## Remarks:

- If  $f$  is continuous at  $x$ , then  $(f(x+) + f(x-))/2 = f(x)$ . So  $f$  equals its Fourier series at “most points.”
- If  $f$  is continuous everywhere, then  $f$  equals its Fourier series everywhere.


A continuous  $2\pi$ -periodic function equals its Fourier series.



A discontinuous  $2\pi$ -periodic piecewise smooth function...



...is *almost* its Fourier series.



# Inner products and orthogonality in $\mathbb{R}^n$

Given vectors  $\mathbf{x} = (x_1, x_2, \dots, x_n)$ ,  $\mathbf{y} = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ , their *inner (or dot) product* is

$$\langle \mathbf{x}, \mathbf{y} \rangle = x_1 y_1 + x_2 y_2 + \dots + x_n y_n.$$

We say that  $\mathbf{x}$  and  $\mathbf{y}$  are *orthogonal* if  $\langle \mathbf{x}, \mathbf{y} \rangle = 0$ .

## Useful facts from linear algebra:

- For any  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}^n$  and  $a, b \in \mathbb{R}$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle \text{ and } \langle a\mathbf{x} + b\mathbf{y}, \mathbf{z} \rangle = a \langle \mathbf{x}, \mathbf{z} \rangle + b \langle \mathbf{y}, \mathbf{z} \rangle.$$

- A set of  $n$  orthogonal vectors in  $\mathbb{R}^n$  forms a basis for  $\mathbb{R}^n$  (an *orthogonal basis*).

Let  $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n$  be orthogonal vectors in  $\mathbb{R}^n$ .

According to the facts above, given  $\mathbf{x} \in \mathbb{R}^n$ , there are (unique) coefficients  $a_1, a_2, \dots, a_n$  so that

$$\mathbf{x} = a_1 \mathbf{b}_1 + a_2 \mathbf{b}_2 + \cdots + a_n \mathbf{b}_n.$$

We can use the inner product to help us compute these coefficients, e.g.

$$\begin{aligned}\langle \mathbf{b}_1, \mathbf{x} \rangle &= \langle \mathbf{b}_1, a_1 \mathbf{b}_1 + a_2 \mathbf{b}_2 + \cdots + a_n \mathbf{b}_n \rangle \\ &= a_1 \langle \mathbf{b}_1, \mathbf{b}_1 \rangle + a_2 \langle \mathbf{b}_1, \mathbf{b}_2 \rangle + \cdots + a_n \langle \mathbf{b}_1, \mathbf{b}_n \rangle \\ &= a_1 \langle \mathbf{b}_1, \mathbf{b}_1 \rangle + 0 + \cdots + 0,\end{aligned}$$

which shows  $a_1 = \langle \mathbf{b}_1, \mathbf{x} \rangle / \langle \mathbf{b}_1, \mathbf{b}_1 \rangle$ .

In general we have the following result.

### Theorem

If  $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n$  is an orthogonal basis of  $\mathbb{R}^n$  and  $\mathbf{x} \in \mathbb{R}^n$ , then

$$\mathbf{x} = a_1\mathbf{b}_1 + a_2\mathbf{b}_2 + \cdots + a_n\mathbf{b}_n$$

where

$$a_i = \frac{\langle \mathbf{b}_i, \mathbf{x} \rangle}{\langle \mathbf{b}_i, \mathbf{b}_i \rangle} \quad (i = 1, 2, \dots, n).$$

### Example

Show that the vectors  $\mathbf{b}_1 = (1, 0, 1)$ ,  $\mathbf{b}_2 = (1, 1, -1)$  and  $\mathbf{b}_3 = (-1, 2, 1)$  form an orthogonal basis for  $\mathbb{R}^3$ , and express  $\mathbf{x} = (1, 2, 3)$  in terms of this basis.

It is easy to see that  $\langle \mathbf{b}_1, \mathbf{b}_2 \rangle = \langle \mathbf{b}_1, \mathbf{b}_3 \rangle = \langle \mathbf{b}_2, \mathbf{b}_3 \rangle = 0$ , e.g.

$$\langle \mathbf{b}_1, \mathbf{b}_2 \rangle = 1 \cdot 1 + 0 \cdot 1 + 1 \cdot (-1) = 0.$$

The theorem tells us that the coordinates of  $\mathbf{x} = (1, 2, 3)$  relative to this basis are

$$\begin{aligned} a_1 &= \frac{\langle \mathbf{b}_1, \mathbf{x} \rangle}{\langle \mathbf{b}_1, \mathbf{b}_1 \rangle} = \frac{1 \cdot 1 + 0 \cdot 2 + 1 \cdot 3}{1 \cdot 1 + 0 \cdot 0 + 1 \cdot 1} = 2, \\ a_2 &= \frac{\langle \mathbf{b}_2, \mathbf{x} \rangle}{\langle \mathbf{b}_2, \mathbf{b}_2 \rangle} = \frac{1 \cdot 1 + 1 \cdot 2 + (-1) \cdot 3}{1 \cdot 1 + 1 \cdot 1 + (-1) \cdot (-1)} = 0, \\ a_3 &= \frac{\langle \mathbf{b}_3, \mathbf{x} \rangle}{\langle \mathbf{b}_3, \mathbf{b}_3 \rangle} = \frac{(-1) \cdot 1 + 2 \cdot 2 + 1 \cdot 3}{(-1) \cdot (-1) + 2 \cdot 2 + 1 \cdot 1} = 1. \end{aligned}$$

That is,

$$\mathbf{x} = 2\mathbf{b}_1 + \mathbf{b}_3.$$

# Inner products of functions

**Goal:** Find an inner product of functions that will allow us to “extract” Fourier coefficients.

**Definition:** Given two functions  $f(x)$  and  $g(x)$ , their *inner product* (on the interval  $[a, b]$ ) is

$$\langle f, g \rangle = \int_a^b f(x)g(x) dx.$$

**Example:** The inner product of  $x$  and  $x^2$  on  $[0, 1]$  is

$$\langle x, x^2 \rangle = \int_0^1 x \cdot x^2 dx = \frac{x^4}{4} \Big|_0^1 = \frac{1}{4}.$$



**Remarks:** If  $f, g, h$  are functions and  $c, d$  are constants, then:

- $\langle f, g \rangle = \langle g, f \rangle$ ;
- $\langle cf + dg, h \rangle = \int_a^b (cf(x) + dg(x))h(x) dx$   
 $= c \int_a^b f(x)h(x) dx + d \int_a^b g(x)h(x) dx$   
 $= c \langle f, h \rangle + d \langle g, h \rangle$ ;
- $\langle f, f \rangle = \int_a^b f(x)^2 dx \geq 0$ ;
- $\langle f, f \rangle = 0$  iff  $f \equiv 0$ ;
- we say  $f$  and  $g$  are *orthogonal on*  $[a, b]$  if  $\langle f, g \rangle = 0$ .

# Examples

1. The functions  $1 - x^2$  and  $x$  are orthogonal on  $[-1, 1]$  since

$$\langle 1 - x^2, x \rangle = \int_{-1}^1 (1 - x^2)x \, dx = \left. \frac{x^2}{2} - \frac{x^4}{4} \right|_{-1}^1 = 0.$$

2. The functions  $\sin x$  and  $\cos x$  are orthogonal on  $[-\pi, \pi]$  since

$$\langle \sin x, \cos x \rangle = \int_{-\pi}^{\pi} \sin x \cos x \, dx = \left. \frac{\sin^2 x}{2} \right|_{-\pi}^{\pi} = 0.$$

3. More generally, for any  $m, n \in \mathbb{N}_0$ , the functions  $\sin(mx)$  and  $\cos(nx)$  are orthogonal on  $[-\pi, \pi]$  since

$$\langle \sin(mx), \cos(nx) \rangle = \int_{-\pi}^{\pi} \underbrace{\sin(mx) \cos(nx)}_{\text{odd}} \, dx = 0.$$

4. For any  $m, n \in \mathbb{N}_0$ , consider  $\cos(mx)$  and  $\cos(nx)$  on  $[-\pi, \pi]$ .

$$\begin{aligned}
 \langle \cos(mx), \cos(nx) \rangle &= \int_{-\pi}^{\pi} \cos(mx) \cos(nx) \, dx \\
 &= \frac{1}{2} \int_{-\pi}^{\pi} \cos((m+n)x) + \cos((m-n)x) \, dx \\
 (m \neq n) \quad &= \frac{1}{2} \left( \frac{\sin((m+n)x)}{m+n} + \frac{\sin((m-n)x)}{m-n} \right) \Big|_{-\pi}^{\pi} \\
 &= 0
 \end{aligned}$$

since  $\sin(k\pi) = 0$  for  $k \in \mathbb{Z}$ . If  $m = n$ , then we have

$$\begin{aligned}
 \langle \cos(mx), \cos(nx) \rangle &= \frac{1}{2} \int_{-\pi}^{\pi} \cos(2mx) + 1 \, dx \\
 (m \neq 0) \quad &= \frac{1}{2} \left( \frac{\sin(2mx)}{2m} + x \right) \Big|_{-\pi}^{\pi} = \pi.
 \end{aligned}$$

Finally, if  $m = n = 0$ , then

$$\langle \cos(mx), \cos(nx) \rangle = \langle 1, 1 \rangle = \int_{-\pi}^{\pi} dx = 2\pi.$$

We conclude that on  $[-\pi, \pi]$  one has

$$\langle \cos(mx), \cos(nx) \rangle = \begin{cases} 0 & \text{if } m \neq n, \\ \pi & \text{if } m = n \neq 0, \\ 2\pi & \text{if } m = n = 0. \end{cases}$$

5. Likewise, on  $[-\pi, \pi]$  one can show that for  $m, n \in \mathbb{N}$

$$\langle \sin(mx), \sin(nx) \rangle = \begin{cases} 0 & \text{if } m \neq n, \\ \pi & \text{if } m = n. \end{cases}$$

# Conclusion

Relative to the inner product

$$\langle f, g \rangle = \int_{-\pi}^{\pi} f(x)g(x) dx,$$

the functions occurring in every Fourier series, namely

$$1, \cos(x), \cos(2x), \cos(3x) \dots, \sin(x), \sin(2x), \sin(3x), \dots$$

form an orthogonal set.

**Moral:** We can use the inner product above to “extract” Fourier coefficients via integration!