

# An Introduction to Partial Differential Equations

Ryan C. Daileda



Trinity University

Partial Differential Equations  
Lecture 1

# Ordinary differential equations (ODEs)

These are equations of the form

$$F(x, y, y', y'', y''', \dots) = 0 \quad (1)$$

where:

- $y = y(x)$  is an (unknown) function of the independent variable  $x$ .
- $y$  is a *solution* of (1) provided the equation holds for *all*  $x$  (in the domain specified).
- The highest derivative occurring in (1) is called the *order* of the equation.

## Some familiar ODEs

You've probably seen the following examples in Calculus II.

1. The solutions of the ODE  $y' = ky$  are  $y = Ce^{kx}$ , for an arbitrary constant  $C$ .
2. The solutions of the ODE  $y'' - y = 0$  are  $y = C_1e^t + C_2e^{-t}$ , for arbitrary constants  $C_1$  and  $C_2$ .
3. The solutions of the ODE  $y'' + y = e^t$  are  $y = C_1 \cos t + C_2 \sin t + \frac{1}{2}e^t$ , for arbitrary constants  $C_1$  and  $C_2$ .

You may want to go back and familiarize yourself with just how these solutions are found.

# The definition of a Partial Differential Equation (PDE)

PDEs are the multivariable analogues of ODEs. As such, they involve *partial derivatives* of an unspecified function.

Specifically, a *Partial Differential Equation* (PDE) has the form

$$F(x_1, x_2, \dots, x_n, u, u_{x_1}, u_{x_2}, \dots, u_{x_n}, \underbrace{\dots\dots\dots}) = 0 \quad (2)$$

higher order partial derivatives of  $u$

where:

- $u = u(x_1, x_2, \dots, x_n)$  is an (unknown) function of the independent variables  $x_1, x_2, \dots, x_n$ .
- $u$  is a *solution* of (2) provided the equation holds for *all*  $x_1, x_2, \dots, x_n$  (in the domain specified).

## Remarks:

Regarding the general PDE

$$F(x_1, x_2, \dots, x_n, u, u_{x_1}, u_{x_2}, \dots, u_{x_n}, \underbrace{\dots\dots\dots}) = 0. \quad (3)$$

higher order partial derivatives of  $u$

1. Recall that  $u_x = \frac{\partial u}{\partial x}$ ,  $u_{xy} = \frac{\partial^2 u}{\partial x \partial y}$ , etc. We will use these notations interchangeably.
2. Although every PDE can be put in the form (3), this is not always necessary.
3. When  $n \leq 4$ , we usually use more familiar independent variables, e.g.  $x, y, z, t$ .
4. The *order* of the PDE (3) is the highest (partial) derivative that explicitly occurs in the equation.

# Examples

1. The function  $u(x, y) = x^2 + y^2$  solves the (first order) PDE  $xu_x + yu_y = 2u$ .
2. The function  $u(x, y) = e^{x-y}$  solves the (first order) PDE  $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} = 0$ .
3. The function  $u(x, y) = x^2 - y^2$  solves the (second order) PDE  $u_{xx} + u_{yy} = 0$ .
4. The function  $u(x, y) = (\sin x)(e^y + e^{-y})$  *also* solves  $u_{xx} + u_{yy} = 0$ .
5. The function  $u(x, y, z) = xyz$  solves the (third order) PDE  $u_{xyz} = 1$ .

## More remarks

1. As with ODEs, checking that a *given* function solves a PDE is straightforward.
2. The hard part is *finding* the solutions to a given PDE.
  - Solution spaces tend to be *infinite dimensional*. It's not usually possible to write down *every* solution.
  - There are no known techniques that will solve all PDEs.

## More remarks

3. However, there are some very powerful techniques that are available for certain classes of PDEs:
  - Method of characteristics
  - Separation of variables, principle of superposition and Fourier series
  - Sturm-Liouville theory
  
4. One can also approximate solutions via numerical methods.
  - Often necessary for extremely complicated problems.
  - Usually studied in other courses, e.g. Heat Transfer.



# Physical phenomena

Many physical phenomena can be effectively modeled via PDEs.

Before we can state them, recall that

$$\nabla = \frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} + \dots$$

is the *gradient operator* and

$$\Delta = \nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \dots$$

is the *Laplacian*.

# Examples

1. **The wave equation:** If  $u(\mathbf{x}, t)$  measures the displacement of an ideal elastic membrane from its equilibrium position, then  $u$  satisfies the (second order) PDE

$$\frac{\partial^2 u}{\partial t^2} = c^2 \Delta u.$$

2. **The heat equation:** If  $u(\mathbf{x}, t)$  gives the temperature in a perfectly thermally conductive medium, then  $u$  satisfies the (second order) PDE

$$\frac{\partial u}{\partial t} = c^2 \Delta u.$$

## Examples

- 3. The transport equation:** If  $u(\mathbf{x}, t)$  is the concentration of a contaminant flowing through a fluid moving with velocity  $\mathbf{v}$ , then  $u$  satisfies the (first order) PDE

$$\frac{\partial u}{\partial t} + \mathbf{v} \cdot \nabla u = 0.$$

- 4. The Laplace equation:** If  $u(\mathbf{x})$  is the steady state temperature in a perfectly thermally conductive medium, then  $u$  satisfies the (second order) PDE

$$\Delta u = 0.$$

## Examples

5. **The (1-D) KdV equation:** If  $u(x, t)$  is the vertical displacement of a flowing shallow fluid, then  $u$  satisfies the (third order) PDE

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + \frac{\partial^3 u}{\partial x^3} = 0.$$


6. **The Schrödinger equation:** If  $u(\mathbf{x}, t)$  is the wave-function of a quantum particle with mass  $\mu$ , subject to a potential  $V(\mathbf{x})$ , then  $u$  satisfies the (second order) PDE

$$i\hbar \frac{\partial u}{\partial t} = -\frac{\hbar^2}{2\mu} \Delta u + V(\mathbf{x})u.$$

# The 1-D transport equation

The set up:

- Consider a fluid flowing with velocity  $v$  through a capillary (of unbounded length) with cross-sectional area  $A$ .
- We introduce a “contaminant” to the fluid, and let  $u(x, t)$  denote its concentration at position  $x$  and time  $t$ .



- At a fixed time  $t$ , the total amount of contaminant between positions  $x_1$  and  $x_2$  is

$$T_1(x_1, x_2, t) = \int_{x_1}^{x_2} u(x, t) \cdot A \, dx. \quad (4)$$

- Similarly, at a fixed position  $x$ , the total amount of contaminant that flows through from time  $t_1$  to  $t_2$  is

$$T_2(t_1, t_2, x) = \int_{t_1}^{t_2} u(x, t) \cdot A \cdot v \, dt. \quad (5)$$

We now “compute the same quantity in two different ways.”

According to (4), the change in the amount of contaminant in the interval  $[x_1, x_2]$  from time  $t_1$  to  $t_2$  is

$$\begin{aligned} T_1(x_1, x_2, t_2) - T_1(x_1, x_2, t_1) &= A \int_{x_1}^{x_2} u(x, t_2) - u(x, t_1) dx \\ &= A \int_{x_1}^{x_2} \int_{t_1}^{t_2} u_t(x, t) dt dx, \quad (6) \end{aligned}$$

where we have used the Fundamental Theorem of Calculus in the final line.

By the same token, (5) tells us the same quantity is also given by

$$\begin{aligned}T_2(t_1, t_2, x_1) - T_2(t_1, t_2, x_2) &= Av \int_{t_1}^{t_2} u(x_1, t) - u(x_2, t) dt \\ &= Av \int_{t_1}^{t_2} \int_{x_2}^{x_1} u_x(x, t) dx dt \\ &= -Av \int_{x_1}^{x_2} \int_{t_1}^{t_2} u_x(x, t) dt dx \quad (7)\end{aligned}$$

where we have used Fubini's theorem to reverse the order of integration (assuming the partial derivatives of the function  $u(x, t)$  are sufficiently smooth).



Since we have simply computed the same quantity in two different ways, (6) and (7) are, in fact, the same:

$$A \int_{x_1}^{x_2} \int_{t_1}^{t_2} u_t(x, t) dt dx = -Av \int_{x_1}^{x_2} \int_{t_1}^{t_2} u_x(x, t) dt dx.$$

Moving everything to one side of the equation yields

$$A \int_{x_1}^{x_2} \int_{t_1}^{t_2} u_t(x, t) + vu_x(x, t) dt dx = 0.$$

# The result

Since  $A > 0$  and  $x_1, x_2, t_1, t_2$  are arbitrary, this can only occur provided

$$u_t(x, t) + vu_x(x, t) = 0$$

for all  $(x, t)$ .

Or, equivalently,

$$\frac{\partial u}{\partial t} + v \frac{\partial u}{\partial x} = 0.$$

This is the **one-dimensional transport equation**.

# Now what?

- Can we even produce *a single* solution to the transport equation?

Yes:  $u(x, t) = x - vt$  works.

- Can we possibly find *every* solution? If so, by what means?

Yes:  $u(x, t) = f(x - vt)$ , where  $f$  is arbitrary.

- We'll answer both of these next time!