

The two dimensional wave equation

R. C. Daileda



Trinity University

Partial Differential Equations
Lecture 11

Vibrating membranes

Goal: Model the motion of an ideal elastic membrane.

Set up: Assume the membrane at rest is a region of the xy -plane and let

$$u(x, y, t) = \begin{cases} \text{vertical deflection of membrane from equilibrium at position } (x, y) \text{ and time } t. \end{cases}$$



For a fixed t , the surface $z = u(x, y, t)$ gives the shape of the membrane at time t .

Under ideal assumptions (e.g. uniform membrane density, uniform tension, no resistance to motion, small deflection, etc.) one can show that u satisfies the *two dimensional wave equation*

$$u_{tt} = c^2 \Delta u = c^2 (u_{xx} + u_{yy}).$$

Rectangular membranes

We assume the membrane lies over the rectangular region $R = [0, a] \times [0, b]$ and has fixed edges.



These facts are expressed by the *boundary conditions*

$$u(0, y, t) = u(a, y, t) = 0, \quad 0 \leq y \leq b, t > 0,$$

$$u(x, 0, t) = u(x, b, t) = 0, \quad 0 \leq x \leq a, t > 0.$$

We must also specify how the membrane is initially deformed and set into motion. This is done via the *initial conditions*

$$\begin{aligned}u(x, y, 0) &= f(x, y), & (x, y) &\in R, \\u_t(x, y, 0) &= g(x, y), & (x, y) &\in R.\end{aligned}$$

New goal: solve the 2-D wave equation subject to the boundary and initial conditions just given.

As usual, we will:

- Use separation of variables to find separated solutions satisfying the homogeneous boundary conditions; and
- Use the principle of superposition to build up a series solution that satisfies the initial conditions as well.

Separation of variables

We seek nontrivial solutions of the form

$$u(x, y, t) = X(x)Y(y)T(t).$$

Plugging this into $u_{tt} = c^2(u_{xx} + u_{yy})$ we get

$$XYT'' = c^2(X''YT + XY''T) \Rightarrow \frac{T''}{c^2T} = \frac{X''}{X} + \frac{Y''}{Y}.$$

Because the two sides are functions of different independent variables, they must be constant:

$$\frac{T''}{c^2T} = A = \frac{X''}{X} + \frac{Y''}{Y} \Rightarrow \begin{cases} T'' - c^2AT = 0, \\ \frac{X''}{X} = -\frac{Y''}{Y} + A. \end{cases}$$

Since the two sides again involve unrelated variables, both are constant:

$$\frac{X''}{X} = B = -\frac{Y''}{Y} + A.$$

Setting $C = A - B$, these equations can be rewritten as

$$X'' - BX = 0, \quad Y'' - CY = 0.$$

The first boundary condition is

$$0 = u(0, y, t) = X(0)Y(y)T(t).$$

Canceling Y and T yields $X(0) = 0$. Likewise, we obtain

$$X(a) = 0, \quad Y(0) = Y(b) = 0.$$

There are no boundary conditions on T .

We have already solved the two boundary value problems for X and Y . The nontrivial solutions are

$$X = X_m(x) = \sin(\mu_m x), \quad \mu_m = \frac{m\pi}{a}, \quad m \in \mathbb{N},$$

$$Y = Y_n(y) = \sin(\nu_n y), \quad \nu_n = \frac{n\pi}{b}, \quad n \in \mathbb{N},$$

with separation constants $B = -\mu_m^2$ and $C = -\nu_n^2$.

Since $T'' - c^2 AT = 0$, and $A = B + C = -(\mu_m^2 + \nu_n^2) < 0$,

$$T = T_{mn}(t) = B_{mn} \cos(\lambda_{mn} t) + B_{mn}^* \sin(\lambda_{mn} t),$$

where

$$\lambda_{mn} = c \sqrt{\mu_m^2 + \nu_n^2} = c\pi \sqrt{\frac{m^2}{a^2} + \frac{n^2}{b^2}}.$$

These are the **characteristic frequencies** of the membrane.

Normal modes

Assembling our results, we find that for any *pair* $m, n \in \mathbb{N}$ we have the *normal mode*

$$\begin{aligned} u_{mn}(x, y, t) &= X_m(x) Y_n(y) T_{mn}(t) \\ &= \sin(\mu_m x) \sin(\nu_n y) (B_{mn} \cos(\lambda_{mn} t) + B_{mn}^* \sin(\lambda_{mn} t)) \\ &= A_{mn} \sin(\mu_m x) \sin(\nu_n y) \cos(\lambda_{mn} t - \phi_{mn}) \end{aligned}$$

Remarks: Note that the normal modes:

- oscillate spatially with frequency $\mu_m/2\pi = m/2a$ in the x -direction,
- oscillate spatially with frequency $\nu_n/2\pi = n/2b$ in the y -direction,
- oscillate temporally with frequency $\lambda_{mn}/2\pi$.
- While $\mu_m/2\pi$ and $\nu_n/2\pi$ are simply multiples of $1/2a$ and $1/2b$, respectively, $\lambda_{mn}/2\pi$ is *not a multiple of any basic frequency*.

Superposition and initial conditions

Superposition gives the general solution

$$u(x, y, t) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sin(\mu_m x) \sin(\nu_n y) (B_{mn} \cos(\lambda_{mn} t) + B_{mn}^* \sin(\lambda_{mn} t)).$$

The initial conditions will determine the coefficients B_{mn} and B_{mn}^* .
Setting $t = 0$ yields

$$f(x, y) = u(x, y, 0) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} B_{mn} \sin\left(\frac{m\pi}{a}x\right) \sin\left(\frac{n\pi}{b}y\right),$$

$$g(x, y) = u_t(x, y, 0) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \lambda_{mn} B_{mn}^* \sin\left(\frac{m\pi}{a}x\right) \sin\left(\frac{n\pi}{b}y\right).$$

These are examples of *double Fourier series*.

Representability

Which functions are given by double Fourier series?

The following result partially answers this first question.

Theorem

If $f(x, y)$ is a C^2 function on the rectangle $[0, a] \times [0, b]$, then

$$f(x, y) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} B_{mn} \sin\left(\frac{m\pi}{a}x\right) \sin\left(\frac{n\pi}{b}y\right),$$

for appropriate B_{mn} .

- To say that $f(x, y)$ is a C^2 function means that f as well as its first and second order partial derivatives are all continuous.
- While not as general as the Fourier representation theorem, this result is sufficient for our applications.

Orthogonality (again!)

How can we compute the coefficients in a double Fourier series?

The following result helps us answer this second question.

Theorem

The functions

$$Z_{mn}(x, y) = \sin\left(\frac{m\pi}{a}x\right) \sin\left(\frac{n\pi}{b}y\right), \quad m, n \in \mathbb{N}$$

are pairwise orthogonal relative to the inner product

$$\langle f, g \rangle = \int_0^a \int_0^b f(x, y)g(x, y) dy dx.$$

This is easily verified using the orthogonality of the functions $\sin(n\pi x/p)$ on the interval $[0, p]$.

Using the usual argument, it follows that if

$$f(x, y) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} B_{mn} \underbrace{\sin\left(\frac{m\pi}{a}x\right) \sin\left(\frac{n\pi}{b}y\right)}_{Z_{mn}},$$

then

$$\begin{aligned} B_{mn} &= \frac{\langle f, Z_{mn} \rangle}{\langle Z_{mn}, Z_{mn} \rangle} = \frac{\int_0^a \int_0^b f(x, y) Z_{mn}(x, y) dy dx}{\int_0^a \int_0^b Z_{mn}(x, y)^2 dy dx} \\ &= \frac{4}{ab} \int_0^a \int_0^b f(x, y) \sin\left(\frac{m\pi}{a}x\right) \sin\left(\frac{n\pi}{b}y\right) dy dx. \end{aligned}$$

So, we can finally write down the complete solution to our original problem.

Conclusion

Theorem

Suppose that $f(x, y)$ and $g(x, y)$ are C^2 functions on the rectangle $[0, a] \times [0, b]$. The solution to the vibrating membrane problem is given by $u(x, y, t) =$

$$\sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sin(\mu_m x) \sin(\nu_n y) (B_{mn} \cos(\lambda_{mn} t) + B_{mn}^* \sin(\lambda_{mn} t))$$

where $\mu_m = \frac{m\pi}{a}$, $\nu_n = \frac{n\pi}{b}$, $\lambda_{mn} = c\sqrt{\mu_m^2 + \nu_n^2}$, and

$$B_{mn} = \frac{4}{ab} \int_0^a \int_0^b f(x, y) \sin(\mu_m x) \sin(\nu_n y) dy dx,$$


$$B_{mn}^* = \frac{4}{ab\lambda_{mn}} \int_0^a \int_0^b g(x, y) \sin(\mu_m x) \sin(\nu_n y) dy dx.$$

Example

A 2×3 rectangular membrane has $c = 6$. If we deform it to have shape given by

$$f(x, y) = xy(2 - x)(3 - y),$$

keep its edges fixed, and release it at $t = 0$, find an expression that gives the shape of the membrane for $t > 0$.



We must compute the coefficients B_{mn} and B_{mn}^* . Since $g(x, y) = 0$ we immediately have

$$B_{mn}^* = 0.$$

We also have

$$\begin{aligned} B_{mn} &= \frac{4}{2 \cdot 3} \int_0^2 \int_0^3 xy(2-x)(3-y) \sin\left(\frac{m\pi}{2}x\right) \sin\left(\frac{n\pi}{3}y\right) dy dx \\ &= \frac{2}{3} \int_0^2 x(2-x) \sin\left(\frac{m\pi}{2}x\right) dx \int_0^3 y(3-y) \sin\left(\frac{n\pi}{3}y\right) dy \\ &= \frac{2}{3} \left(\frac{16(1 + (-1)^{m+1})}{\pi^3 m^3} \right) \left(\frac{54(1 + (-1)^{n+1})}{\pi^3 n^3} \right) \\ &= \frac{576}{\pi^6} \frac{(1 + (-1)^{m+1})(1 + (-1)^{n+1})}{m^3 n^3}. \end{aligned}$$

The coefficients λ_{mn} are given by

$$\lambda_{mn} = c \sqrt{\mu_n^2 + \nu_n^2} = 6\pi \sqrt{\frac{m^2}{4} + \frac{n^2}{9}} = \pi \sqrt{9m^2 + 4n^2}.$$

Assembling all of these pieces yields

$$u(x, y, t) = \frac{576}{\pi^6} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \left(\frac{(1 + (-1)^{m+1})(1 + (-1)^{n+1})}{m^3 n^3} \sin\left(\frac{m\pi}{2}x\right) \right. \\ \left. \times \sin\left(\frac{n\pi}{3}y\right) \cos\left(\pi \sqrt{9m^2 + 4n^2} t\right) \right).$$

Example

Suppose in the previous example we also impose an initial velocity given by $g(x, y) = 8 \sin 2\pi x$. Find an expression that gives the shape of the membrane for $t > 0$.

Since we have the same initial shape, B_{mn} don't change. We only need to find B_{mn}^* and add the appropriate terms to the previous solution.

Using λ_{mn} computed above, we have

$$\begin{aligned} B_{mn}^* &= \frac{4}{2 \cdot 3\pi\sqrt{9m^2 + 4n^2}} \int_0^2 \int_0^3 8 \sin(2\pi x) \sin\left(\frac{m\pi}{2}x\right) \sin\left(\frac{n\pi}{3}y\right) dy dx \\ &= \frac{16}{3\pi\sqrt{9m^2 + 4n^2}} \int_0^2 \sin(2\pi x) \sin\left(\frac{m\pi}{2}x\right) dx \int_0^3 \sin\left(\frac{n\pi}{3}y\right) dy. \end{aligned}$$

The first integral is zero unless $m = 4$, i.e. $B_{mn}^* = 0$ for $m \neq 4$.

Evaluating the second integral, we have

$$B_{4n}^* = \frac{8}{3\pi\sqrt{36+n^2}} \frac{3(1+(-1)^{n+1})}{n\pi} = \frac{8(1+(-1)^{n+1})}{\pi^2 n\sqrt{36+n^2}}.$$

So the velocity dependent term of the solution is

$$\begin{aligned} u_2(x, y, t) &= \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} B_{mn}^* \sin(\mu_m x) \sin(\nu_n y) \sin(\lambda_{mnt}) \\ &= \frac{8 \sin(2\pi x)}{\pi^2} \sum_{n=1}^{\infty} \frac{1+(-1)^{n+1}}{n\sqrt{36+n^2}} \sin\left(\frac{n\pi}{3} y\right) \sin\left(2\pi\sqrt{36+n^2} t\right). \end{aligned}$$

If we let $u_1(x, y, t)$ denote the solution to the first example, the complete solution here is

$$u(x, y, t) = u_1(x, y, t) + u_2(x, y, t).$$